# ASTR 610 Theory of Galaxy Formation

Lecture 19: Elliptical Galaxies

FRANK VAN DEN BOSCH Yale University, Spring 2024



### **Structure & Formation of Ellitpical Galaxies**

In this lecture we discuss the structure and formation of elliptical galaxies. After a very brief overview of some of the main observational properties of ellipticals, we discuss two `competing' pictures for the formation of ellipticals.



### **Observational Facts**

**Surface Brightness Profiles** 

Elliptical galaxies have surface brightness profiles that are well described by a Sersic profile:

$$I(R) = I_0 \exp\left[-\beta_n \left(\frac{R}{R_e}\right)^{1/n}\right]$$

 $eta_n\simeq 2n-0.324$  follows from definition of Re

$$L = 2\pi \int_0^\infty I(R) R \, \mathrm{d}R = \frac{2\pi n \Gamma(2n)}{(\beta_n)^{2n}} I_0 R_e^2$$

n: Sersic index R<sub>e</sub>: effective radius that encloses half of the total light

For n=4 this is known as the `de Vaucouleurs' profile, while n=1 corresponds to an exponential profile



Typically, brighter ellipticals have a larger Sersic index Faintest ellipticals (dwarf spheroidals) have n~1, corresponding to an exponential profile

ASTR 610:Theory of Galaxy Formation

### **Observational Facts**

#### Surface Brightness Profiles



Brighter ellipticals are larger and have higher surface brightness But, trends are not continuous: for  $M_B < -20.5$  surface brightess decreases with increasing luminosity, while there is little to no magnitude-size trend for dwarfs...

### **Isophotal Shapes**

The isophotes of elliptical galaxies are elliptical, but not perfectly so. They show deviations that are typically either 'disky' or 'boxy'



$$\Delta \phi \equiv R_{\rm iso}(\phi) - R_{\rm ell}(\phi)$$
$$= \sum_{n=3}^{\infty} [a_n \cos(n\phi) + b_n \sin(n\phi)]$$

disky and boxy correspond to a<sub>4</sub>>0 and a<sub>4</sub><0, respectively

#### Brighter ellipticals are more likely boxy (disky fraction decreases with luminosity)





#### ASTR 610: Theory of Galaxy Formation

# Isophotal Twist



The presence of isophote twists among (bright/boxy) ellipticals is often taken as evidence that, as a class, they must be triaxial. Some ellipticals reveal isophotal twists, with direction of major axis of isophote changing with isophotal level

Most of these ellipticals are boxy

The simplest explanation is that (these) elliptical galaxies are triaxial (rather than oblate/prolate), and have their intrinsic axis ratios change with radius.

Such a system in projection will reveal isophote twist



ASTR 610: Theory of Galaxy Formation

### **The Nuclei of Elliptical Galaxies**

High resolution imaging with the HST revealed that the central regions of ellipticals reveal a dichotomy in their central surface brightness profile; 'cusps' vs 'cores'



Typically cored ellipticals are bright ( $M_B \leq -20.5$ ) and boxy, while cuspy ellipticals are fainter and disky.

Whether this is a true `dichotomy' or not is still debated...

Nuclei of elliptical galaxies also often harbour small (few 100pc) disks of gas/dust and/or stars

### The Nuclear Dust Disk of NGC 4261



Dust disk ~100 pc size, oriented perpendicular to radio jets; is this the material that feeds the accretion disk surrounding the SMBH at the center?

#### ASTR 610: Theory of Galaxy Formation

# **Creating Cores with Scouring SMBH Binaries**



Milosavljevic & Merritt, 2001, ApJ, 563, 34

#### Rule of thumb Mej ~ 0.5 (M<sub> $\bullet$ ,1</sub> + M<sub> $\bullet$ ,2</sub>) In(a<sub>h</sub>/a<sub>gr</sub>)

- a<sub>h</sub> = semi-major axis of SMBH binary when binary first becomes hard
- a<sub>h</sub> = semi-major axis of SMBH binary when gravitational radiation starts to dominate

Cores can be created due to scouring by a SMBH binary. Dynamical friction acting on the SMBHs tightens the binary, and transfers momentum to the cusp stores, thereby creating a core. This process becomes inefficient once gravitational wave radiation becomes important..



#### ASTR 610: Theory of Galaxy Formation

#### The Nuclear Stellar Disk of NGC 4342



#### ASTR 610:Theory of Galaxy Formation

### Kinematics

The observed spectrum of an elliptical galaxy is a convolution of the template spectrum, which is the luminosity weighted spectrum of all the various stars along the line-of-sight (LOS) and a broadening function, which is a combination of an instrumental broadening function an the line-of-sight velocity distribution (LOSVD)

A typical functional form for the LOSVD is a simple Gaussian.

$$\mathcal{L}(v) = \frac{1}{\sqrt{2\pi\sigma}} e^{-\frac{1}{2}w^2} \qquad w = (v - V)/\sigma$$

However, the LOSVD is generally not Gaussian and is has become standard practice to adopt a Gauss-Hermite series

$$\mathcal{L}(v) = \frac{1}{\sqrt{2\pi\sigma}} e^{-\frac{1}{2}w^2} \left[ 1 + \sum_{j=3}^N h_j H_j(w) \right]$$



van der Marel & Franx 1993

Typically one truncates the series at N=4, such that LOSVD is described by four parameters: V,  $\sigma$ , h<sub>3</sub>, h<sub>4</sub>

related to skewness

related to kurtosis

© Frank van den Bosch, Yale University

ASTR 610: Theory of Galaxy Formation

### **Kinematics**



The h<sub>4</sub> Gauss-Hermite moment is especially powerful as it is sensitive to the orbital distribution of the galaxy, and can therefore be used to break the mass-anisotropy degeneracy that hampers kinematic models.

#### ASTR 610: Theory of Galaxy Formation

### **Kinematics**



Disky ellipticals typically reveal strong rotation along major axis, consistent with them being `oblate rotators' (oblate in shape, with flattening due to rotation)

Boxy ellipticals reveal little rotation, and occasionaly rotation along the minor axis. Latter is a clear sign that (boxy) ellipticals are triaxial

Emsellem

### **Orbital Families in Triaxial Potentials**

box orbit



short-axis tube orbit

outer long-axis tube orbit

In triaxial potentials, there are four families of regular orbits. Box orbit have no net angular momentum; the orbit comes arbitrarily close to the centre. Tube orbits, on the other hand, have an angular momentum barrier

We now examine how the structure of elliptical galaxies relates to their kinematics.

The dynamics of (elliptical) galaxies are governed by the CBE: df/dt = 0

Mutiplying the CBE with velocity, and integrating over velocity-space yields the Jeans equations (which are momentum equations)

$$\frac{\partial(\rho\langle v_j\rangle)}{\partial t} + \frac{\partial(\rho\langle v_iv_j\rangle)}{\partial x_i} + \rho\frac{\partial\Phi}{\partial x_j} = 0$$

Multiplying all terms with  $x_k$  and integrating over all of configuration space yields

$$\frac{\partial}{\partial t} \int \rho x_k \langle v_j \rangle \, \mathrm{d}^3 \vec{x} = -\int x_k \frac{\partial (\rho \langle v_i v_j \rangle)}{\partial x_i} \, \mathrm{d}^3 \vec{x} - \int \rho x_k \frac{\partial \Phi}{\partial x_j} \mathrm{d}^3 \vec{x}$$

Using integration by parts, the first terms on the rhs can be written as

$$\int x_k \frac{\partial (\rho \langle v_i v_j \rangle)}{\partial x_i} \, \mathrm{d}^3 \vec{x} = -\int \rho \langle v_k v_j \rangle \mathrm{d}^3 \vec{x} \equiv -2\mathcal{K}_{kj}$$

where we have defined the kinetic energy tensor,  $\mathcal{K}_{kj}$ 

#### ASTR 610: Theory of Galaxy Formation

# Centrifugal Support vs. Pressure Support

We split the kinetic energy tensor into contributions from ordered and random motions:

$$\mathcal{T}_{ij} \equiv \frac{1}{2} \int \rho \langle v_i \rangle \langle v_j \rangle d^3 \vec{x}$$
$$\Pi_{ij} \equiv \int \rho \sigma_{ij}^2 d^3 \vec{x}$$

In addition to the kinetic energy tensor, we also define the potential energy tensor:

$$\mathcal{W}_{ij} \equiv -\int \rho x_i \frac{\partial \Phi}{\partial x_j} \,\mathrm{d}^3 \vec{x}$$

Combining the above, and using that both  $\mathcal{K}$  and  $\mathcal{W}$  are symmetric, we have that

$$\frac{1}{2}\frac{\mathrm{d}}{\mathrm{d}t}\int\rho\left[x_k\langle v_j\rangle+x_j\langle v_k\rangle\right]\,\mathrm{d}^3\vec{x}=2\mathcal{K}_{jk}+\mathcal{W}_{jk}$$

ASTR 610:Theory of Galaxy Formation

Finally, we define the moment of inertia tensor

$$\mathcal{I}_{ij} \equiv \int \rho x_i x_j \mathrm{d}^3 \vec{x}$$

Differentiating wrt time and using the continuity equation (i.e., zeroth moment of CBE):

$$\frac{\mathrm{d}\mathcal{I}_{jk}}{\mathrm{d}t} = \int \frac{\partial\rho}{\partial t} x_j x_k \,\mathrm{d}^3 \vec{x} = -\int \frac{\partial\rho\langle v_i \rangle}{\partial x_i} x_j x_k \mathrm{d}^3 \vec{x} = \int \rho \left[ x_j \langle v_k \rangle + x_k \langle v_j \rangle \right] \,\mathrm{d}^3 \vec{x}$$

students: try this at home

so that

$$\frac{1}{2}\frac{\mathrm{d}}{\mathrm{d}t}\int\rho\left[x_{j}\langle v_{k}\rangle+x_{k}\langle v_{j}\rangle\right]\,\mathrm{d}^{3}\vec{x}=\frac{1}{2}\frac{\mathrm{d}^{2}\mathcal{I}_{jk}}{\mathrm{d}t^{2}}$$

which allows us to write the Tensor Virial Theorem as

$$\frac{1}{2} \frac{\mathrm{d}^2 \mathcal{I}_{jk}}{\mathrm{d}t^2} = 2\mathcal{T}_{jk} + \Pi_{jk} + \mathcal{W}_{jk}$$

which relates the gross kinematic and structural properties of gravitational systems

If the system is in a steady state, the moment of inertia tensor is stationary, and the tensor virial theorem reduces to

 $2\mathcal{K}_{jk} + \mathcal{W}_{jk} = 0$ 

The common (scalar) virial theorem (2K+W=0) is simply the trace of this tensor equation.

We now use this tensor virial theorem, to relate the flattening of an elliptical to its kinematics. Consider an oblate system with it's symmetry axis along the z-direction. Because of symmetry considerations we have that

$$\langle v_R \rangle = \langle v_z \rangle = 0$$
  $\langle v_R v_\phi \rangle = \langle v_z v_\phi \rangle = 0$ 

If we write  $\langle v_x \rangle = \langle v_\phi \rangle \sin \phi$  and  $\langle v_y \rangle = \langle v_\phi \rangle \cos \phi$  then we obtain

$$\mathcal{T}_{xy} = \frac{1}{2} \int \rho \langle v_x \rangle \langle v_y \rangle \,\mathrm{d}^3 \vec{x} = \frac{1}{2} \int_0^{2\pi} \mathrm{d}\phi \sin\phi \,\cos\phi \int_0^{\infty} \mathrm{d}R \int_{-\infty}^{+\infty} \mathrm{d}z \,\rho(R,z) \,\langle v_\phi \rangle^2(R,z) = 0$$

A similar analysis shows that all other non-diagonal elements of  $\mathcal{T}$ ,  $\Pi$  and  $\mathcal{W}$  have to be zero In addition, because of symmetry considerations we must also have that  $\mathcal{T}_{xx} = \mathcal{T}_{yy}$ , and similar for  $\Pi$  and  $\mathcal{W}$ 

Given these symmetries, the only independent, non-trivial virial equations are

$$2\mathcal{T}_{xx} + \Pi_{xx} + \mathcal{W}_{xx} = 0 \qquad \qquad 2\mathcal{T}_{zz} + \Pi_{zz} + \mathcal{W}_{zz} = 0$$

If the only streaming motion is rotation about the z-axis, then  $T_{zz} = 0$  and

$$2\mathcal{T}_{xx} = \frac{1}{2} \int \rho \langle v_{\phi} \rangle^2 \mathrm{d}^3 \vec{x} = \frac{1}{2} M v_0^2$$

where  $v_0$  is the mass-weighted rotation velocity. Similarly we can write

$$\Pi_{xx} = M\sigma_0^2; \qquad \Pi_{zz} = (1-\delta)M\sigma_0^2$$

where  $\sigma_0^2 \equiv (1/M) \int \rho \sigma_{xx}^2 d^3 \vec{x}$  is the mass-weighted velocity dispersion along los, and  $\delta \equiv 1 - \prod_{zz} / \prod_{xx} < 1$  is a measure of the anisotropy of the velocity dispersion

Taking the ratio between the two non-trivial virial equations above then yields

$$\frac{\mathcal{W}_{xx}}{\mathcal{W}_{zz}} = \frac{1}{1-\delta} \left( 1 + \frac{1}{2} \frac{v_0^2}{\sigma_0^2} \right)$$

#### ASTR 610: Theory of Galaxy Formation

As shown by Roberts (1962), for systems stratified on similar coaxial oblate ellipsoids, the ratio  $W_{xx}/W_{zz}$  depends only on the ellipticity  $\varepsilon$ 

$$\frac{\mathcal{W}_{xx}}{\mathcal{W}_{zz}} = \frac{1}{1-\delta} \left( 1 + \frac{1}{2} \frac{v_0^2}{\sigma_0^2} \right)$$

The above expression therefore makes it clear that a stellar system can be flattened either by rotation, or by anisotropic velocity dispersion (i.e.,  $\delta > 0$ )

It is customary to identify  $\sigma_0$  with  $\bar{\sigma}$ , the mean velocity dispersion interior to half the effective radius, and  $v_0$  with  $4v_m/\pi$ , with  $v_m$  the maximum rotation velocity (Binney 2005)



Solid lines are for edge-on system; dashed lines show impact of projection for decreasing inclination angle.

For isotropic case, to good approximation we have

$$\frac{v_{\rm m}}{\bar{\sigma}} \approx \sqrt{\frac{\varepsilon}{1-\varepsilon}}$$

ASTR 610:Theory of Galaxy Formation

### Fast vs. Slow Rotators





#### One can split ellipticals in two kinematic classes:

Fast rotators; kinematics consistent with oblate rotators, shape is flattened by rotation
Slow rotators; very little rotation; shape is due to anisotropic pressure support

Typically, slow rotators are more massive, and are often boxy. Fast rotators are disky ellipticals or S0s, and are often less luminous.

$$\lambda_R = \frac{\langle R \, |V| \rangle}{\langle R \, \sqrt{V^2 + \sigma^2} \rangle}$$

a modern replacement for 
$$v_{
m m}/ar{\sigma}$$

ASTR 610:Theory of Galaxy Formation

Emsellem et al. 2011, MNRAS, 414, 888

### The Dichotomy among Ellipticals



Disky ellipticals are consistent with being more bulge-dominated versions of S0 galaxies.

#### ASTR 610:Theory of Galaxy Formation

### **The Fundamental Plane**



Similar to the TF-relation for disk galaxies, ellipticals reveal a scaling relation between luminosity and velocity dispersion, known as the Faber-Jackson (FJ) relation:

However, unlike the TF, the scatter in FJ \*is\* correlated with size, giving rise to a three-parameter Fundamental Plane relation.



#### ASTR 610: Theory of Galaxy Formation

### **The Fundamental Plane**

The FP-relation is generally written in the form

 $\log R_{\rm e} = a \log \sigma_0 + b \log \langle I \rangle_{\rm e} + \text{cst}$ 

Best-fit parameters are  $a \sim 1.2$  to 1.5 (depending on waveband), and  $b \sim -0.8$ 

The FP-relation is usually interpreted in terms of the Virial Theorem

$$\frac{GM}{\langle R\rangle} = \langle v^2 \rangle$$

 $\langle R \rangle$  = average radius, such that lhs is abs. value of mean potential energy per unit mass  $\langle v^2 \rangle$  = average rms velocity, such that half that value is mean kinetic energy per unit mass

If ellipticals are homologous (i.e.  $\kappa_R$  and  $\kappa_V$  constant), and the mass-to-light ratio is constant, then the Virial Theorem predicts a FP-relation with a=2 and b=-1

The deviation from this prediction is called the 'tilt' of the fundamental plane, and reflects that ellipticals as a class are not homologous, and/or that  $(M/L) \propto L^{\alpha} \langle I \rangle_{e}^{\beta}$  with  $(\alpha, \beta) \neq 0$ .

although still debated, the latter option seems to explain most of the tilt.

#### ASTR 610: Theory of Galaxy Formation

### **The Fundamental Plane**

As originally proposed by Bender+92, it is useful to use an orthogonal combination of the three observables that enter the FP-relation, which facilitates interpretation.

 $\kappa_{1} \equiv (\log \sigma_{0}^{2} + \log R_{e})/\sqrt{2}$   $\kappa_{2} \equiv (\log \sigma_{0}^{2} + 2 \log \langle I \rangle_{e} - \log R_{e})/\sqrt{6}$  $\kappa_{3} \equiv (\log \sigma_{0}^{2} - \log \langle I \rangle_{e} - \log R_{e})/\sqrt{3}$ 

 $\kappa_1 \propto \log(\sigma_0^2 R_{\rm e}) \propto \log M$  $\kappa_3 \propto \log(\sigma_0^2 R_{\rm e} / \langle I \rangle_{\rm e} R_{\rm e}^2) \propto \log(M/L)$ 

In this ' $\kappa$ -space', the  $\kappa_1$ - $\kappa_2$  projection is very close to a face-on projection of the FP, while the  $\kappa_1$ - $\kappa_3$  projection shows the FP nearly edge-on. In fact, if ellipticals are homologous, and (M/L)  $\sim$  M<sup> $\gamma$ </sup>, the virial theorem implies that  $\kappa_3 = \sqrt{2/3}\gamma\kappa_1 + \text{cst}$ 



ASTR 610:Theory of Galaxy Formation

### **Formation of Ellitpical Galaxies**

One 'obvious' scenario for why some galaxies are ellipticals and others are spirals is to assume this is governed by angular momentum....





ASTR 610: Theory of Galaxy Formation

© Frank van den Bosch, Yale University